

CANT 2022: Zoom Conference

Twentieth Annual Workshop on
Combinatorial and Additive Number Theory
CUNY Graduate Center
May 24 - 27, 2022

Abstracts (as of May 10, 2022)

- (1) **Emma Bailey**, CUNY Graduate Center

Email: ebailey@gc.cuny.edu

Title: Large deviations of Selberg's Central Limit Theorem

Abstract: Selberg's celebrated central limit theorem shows that $\log \zeta(1/2 + it)$ at a typical point t at height T behaves like a complex, centered Gaussian random variable with variance $\log \log T$. This talk will present recent results showing that the Gaussian decay persists in the large deviation regime, at a level on the order of the variance, improving on the best known bounds in that range. Time permitting, we will also present various applications, including on the maximum of the zeta function in short intervals.

This work is joint with Louis-Pierre Arguin.

- (2) **Benjamin Baily**, Williams College, and

Henry Fleischmann, University of Michigan

Email: bmb2@williams.edu and henryfl@umich.edu

Title: Large sets are sumsets

Abstract: Let $[n] := \{0, 1, 2, \dots, n\}$. Intuitively, all large subsets of $[n]$ have additive structure, and Roth famously made this precise by finding constants $c, N > 0$ such that for $n \geq N$, any subset of $[n]$ containing more than $\frac{cn}{\log \log n}$ elements must contain an arithmetic progression of length 3. We establish a different interpretation of the intuition by finding explicit constants $\alpha = \frac{1}{\log 2}$ and $\beta = \frac{1}{\log 1.325}$ such that, for sufficiently large n , we have:

- (i) any subset of $[n]$ with more than $n - \alpha \log n$ elements has a nontrivial decomposition as the sum of two sets, and
- (ii) there exists a subset of $[n]$ of size $n - \beta \log n$ at least that has no such decomposition.

We also prove, using these methods, a higher-dimensional analogue of results (i) and (ii). Notably, our threshold at which structure appears is far higher than Roth's.

This work was joint with Justine Dell, Sophia Dever, Adam Dionne, Faye Jackson, Leo Goldmakher, Gal Gross, Steven J. Miller, Ethan Pesikoff, Huy Tuan Pham, Luke Reifenberg, and Vidya Venkatesh.

- (3) **Gautami Bhowmik**, Université de Lille
 Email: gautami.bhowmik@univ-lille.fr
 Title: Siegel zeros under Goldbach conjectures
 Abstract: A Landau-Siegel zero is a possible though unwelcome counterexample to the Generalised Riemann Hypothesis. Proving its absence unconditionally is clearly a difficult problem. We will discuss some results by assuming plausible conjectures on the Goldbach problem: the Hardy-Littlewood one (1923), a weak form due to Fei (2016), and a weaker form that we studied more recently (Bhowmik-Halupczok, in: Proceedings of CANT 2019 and 2020). Continuing on these lines, Friedlander-Goldston-Iwaniec-Suriajaya (2022) showed that the assumption of Fei's conjecture is enough to disprove the existence of Siegel zeros.
- (4) **Jörg Brüdern**, Universität Göttingen, Germany,
 Email: joerg.bruedern@mathematik.uni-goettingen.de
 Title: Bracketed ternary additive problems
 Abstract: The ternary additive problems of Waring's type (that is, sums of three potentially unlike powers) have attracted many workers in the additive theory of numbers. In this talk, we discuss several variants that involve brackets (that is, the integer part of certain monomials).
- (5) **Jin-Hui Fang**, Nanjing University of Information Science and Technology
 Email: fangjinhui1114@163.com
 Title: Representation functions avoiding integers with density zero
 Abstract: For a nonempty set A of integers and any integer n , denote $r_A(n)$ by the number of representations of n of the form $n = a + a'$, where $a \leq a'$ and $a, a' \in A$ and $d_A(n)$ by the number of pairs (a, a') with $a, a' \in A$ such that $n = a - a'$. In 2008, Nathanson considered the representation function with infinitely many zeros. Following Nathanson's work, we proved that, for any set T of integers with density zero, there exists a sequence A of integers such that $r_A(n) = 1$ for all integers $n \notin T$ and $r_A(n) = 0$ for all integers $n \in T$, and $d_A(n) = 1$ for all positive integers n . We will also present our recent results on representation functions.
- (6) **Leonid Fel**, Technion – Israel Institute of Technology, Israel
 Email: lfel@cv.technion.ac.il
 Title: Commutative monoid of self-dual symmetric polynomials
 Abstract: We consider a set $\mathfrak{RS}(\lambda, S_n)$ of self- and skew-reciprocal polynomials in λ , of degree mn , where $m \in \mathbb{Z}_{\geq}$, $n \in \mathbb{Z}_{>}$, based on polynomial invariants $I_{n,r}(\mathbf{x}^n)$ of symmetric group S_n , acting on the Euclidean space \mathbb{E}^n over the field of real numbers \mathbb{R} , where $\mathbf{x}^n = \{x_1, \dots, x_n\} \in \mathbb{E}^n$. We prove that $\mathfrak{RS}(\lambda, S_n)$ exhibits a commutative monoid under multiplication. Real solutions $\lambda(\mathbf{x}^n)$ of skew-reciprocal equations have many remarkable properties: a homogeneity of the 1st order, a duality under inversion of variables $x_i \rightarrow x_i^{-1}$ and function $\lambda \rightarrow \lambda^{-1}$, a monotony of $\lambda(\mathbf{x}^n)$ with respect to every x_i and others. We find the bounds of $\lambda(\mathbf{x}^n)$ which are given by arithmetic and harmonic means of the set $\{x_1, \dots, x_n\}$.

- (7) **Henry Fleischmann**, University of Michigan, and
Ethan Pesikoff, Yale University

Email: henryfl@umich.edu and ethan.pesikoff@yale.edu

Title: Angle variants of the Erdős distinct distance problem

Abstract: The Erdős distinct distance problem is a ubiquitous problem in discrete geometry. Less well known is Erdős' distinct angle problem, the problem of finding the minimum number of distinct angles between n non-collinear points in the plane. We provide new upper and lower bounds on a broad class of distinct angle problems. We show that the number of distinct angles formed by n points in general position is $O(n^{\log_2(7)})$, providing the first non-trivial bound for this quantity. We introduce several new asymptotically optimal configurations. We also show, via a probabilistic argument, that the minimum size of a maximal subset of n points in general position admitting only unique angles is $\Omega(n^{1/5})$ and $O(n^{\log_2(7)/3})$. These are the best known bounds on this quantity.

This work is joint with Hongyi Hu, Faye Jackson, Steven J. Miller, Evinur Palsson, and Charles Wolf.

- (8) **Mikhail Gabdullin**, Steklov Mathematical Institute, Moscow, Russia

Email: gabdullin.mikhail@yandex.ru

Title: A conjecture of Cilleruelo and Cordoba and divisors in a short interval

Abstract: Let $E(A) = \#\{(a_1, a_2, a_3, a_4) \in A^4 : a_1 + a_2 = a_3 + a_4\}$ denote the additive energy of a set $A \subset \mathbf{N}$, and let $\mathbb{T} = \mathbf{R}/\mathbf{Z}$ and $\|f\|_4 = (\int_{\mathbb{T}} |f(t)|^4 dt)^{1/4}$. It is well-known that

$$E(\{n^2 : n \leq N\}) = \left\| \sum_{n \leq N} e^{2\pi i n^2 x} \right\|_4^4 \asymp N^2 \log N,$$

while we trivially have $E(A) \geq |A|^2$. In 1992, J. Cilleruelo and A. Cordoba proved that $E(\{n^2 : N \leq n \leq N + N^\gamma\}) \asymp N^{2\gamma}$ for any $\gamma \in (0, 1)$, and conjectured a much more general bound (again, for any $\gamma \in (0, 1)$)

$$\left\| \sum_{N \leq n \leq N + N^\gamma} a_n e^{2\pi i n^2 x} \right\|_4 \leq C(\gamma) \left(\sum_{N \leq n \leq N + N^\gamma} |a_n|^2 \right)^{1/2}.$$

While this bound is easy to prove for $\gamma \leq 1/2$, it seems to be open for any $\gamma > 1/2$. We prove this for all $\gamma < \frac{\sqrt{5}-1}{2} = 0.618\dots$ and present a connection between this problem and a conjecture of I. Ruzsa: for any $\epsilon > 0$ there exists $C(\epsilon) > 0$ such that any positive integer N has at most $C(\epsilon)$ divisors in the interval $[N^{1/2}, N^{1/2} + N^{1/2-\epsilon}]$.

- (9) **Krystian Gajdzica**, Jagiellonian University, Kraków, Poland
 Email: krystian.gajdzica@im.uj.edu.pl
 Title: Some inequalities for the multicolor restricted partition function $p_{\mathcal{A}}(n, k)$
 Abstract: For a non-decreasing sequence of positive integers $\mathcal{A} = (a_i)_{i=1}^{\infty}$ and a fixed integer $k \geq 1$, the multicolor restricted partition function $p_{\mathcal{A}}(n, k)$ counts the number of partitions of n with parts in the multiset $\{a_1, a_2, \dots, a_k\}$. The talk is devoted to some multiplicative inequalities related to $p_{\mathcal{A}}(n, k)$. Among other things, we will examine: the Bessenrodt-Ono inequality for $p_{\mathcal{A}}(n, k)$, the log-concavity of the sequence $(p_{\mathcal{A}}(n, k))_{n=1}^{\infty}$, the higher order Turán property and other similar phenomena.
- (10) **Heidi Goodson**, Brooklyn College (CUNY)
 Email: Heidi.Goodson@brooklyn.cuny.edu
 Title: Sato-Tate groups in dimension greater than 3
 Abstract: The focus of this talk is on Sato-Tate groups of abelian varieties – compact groups predicted to determine the limiting distributions of local zeta functions. In recent years, complete classifications of Sato-Tate groups in dimensions 1, 2, and 3 have been given, but there are obstacles to providing classifications in higher dimensions. In this talk, I will describe my recent work on families of higher dimensional Jacobian varieties.
 This work is partly joint with Melissa Emory.
- (11) **Rachel Greenfeld**, UCLA
 Email: greenfeld@ucla.edu
 Title: Translational tilings
 Abstract: Translational tiling is a covering of a space using translated copies of some building blocks, called the tiles, without any positive measure overlaps. Which are the possible ways that a space can be tiled? In the talk, we will discuss the study of this question as well as its applications, and report on recent progress, joint with Terence Tao.
- (12) **Li Guo**, Rutgers University - Newark
 Email: liguo@rutgers.edu
 Title: Renormalization of quasisymmetric functions
 Abstract: The algebra of quasisymmetric functions (QSym) has played a central role in multiple zeta values and a large class of combinatorial algebraic structures related to symmetric functions. A natural linear basis of QSym is the set of monomial quasisymmetric functions defined by compositions, that is, vectors of positive integers. Extending such a definition for weak compositions, that is, vectors of nonnegative integers, leads to divergent expressions. This phenomenon is closely related to the divergency of multiple zeta values with nonpositive integer arguments.

We apply the method of renormalization in the spirit of Connes and Kreimer to address the divergency, and realize weak composition quasisymmetric functions as power series. The resulting Hopf algebra has the Hopf algebra of quasisymmetric functions as both a Hopf subalgebra and a Hopf quotient algebra.

This is joint work with Houyi Yu and Bin Zhang.

- (13) **Shruti Hegde**, Ramakrishna Mission Vivekananda Educational and Research Institute, India

Email: shrutihg2112@gmail.com

Title: Weighted zero-sum constants and inverse results

Abstract: A weighted generalization of classical zero-sum constants was introduced by Adhikari *et al.* in 2006 and has been an active area of research since then. In the last fifteen years, weighted zero-sum constants for \mathbb{Z}_n with several interesting weight sets have been found. In this talk, we take up the problem of determining the exact values and providing bounds of the weighted Davenport constant of \mathbb{Z}_n with some new weight sets.

Next, we consider a weighted generalization of the *the Erdős-Ginzburg-Ziv constant*. Let G be a finite abelian group with $\exp(G) = n$. For a positive integer k and a non-empty subset A of $[1, n - 1]$, the arithmetical invariant $s_{kn,A}(G)$ is defined to be the least positive integer t such that any sequence S of t elements in G has an A -weighted zero-sum subsequence of length kn . We give the exact value of $s_{kq,A}(G)$, for integers $k \geq 2$ and $A = \{1, 2\}$, where G is an abelian p -group with $\text{rank}(G) \leq 4$, p is an odd prime and $\exp(G) = q$. Our method consists of a modification of a polynomial method of Rónyai.

Lastly, we consider the questions regarding inverse problems for the weighted zero-sum constants of \mathbb{Z}_n . An inverse problem is the problem of characterizing all the weighted *zero-sum free sequences* over \mathbb{Z}_n of specific lengths for the particular weight sets under consideration.

This work was joint with Sukumar Das Adhikari and partly with Md Ibrahim Molla and Subha Sarkar.

- (14) **Norbert Hegyvari**, Eötvös Loránd University and Alfréd Rényi Institute of Mathematics, Hungary

Email: hegyvari@renyi.hu

Title: Boolean functions defined on pseudo-recursive sequences

Abstract: We define Boolean functions on hypergraphs with edges having big intersections, and an opposite situation, hypergraphs which are thinly intersecting induced by pseudo-recursive sequences. A sequence X is said to be *pseudo-recursive sequence* if the identity $x_{n+1} = m_{n+1}x_n + b_{j_{n+1}}$ holds, where $b_{j_{n+1}} \in \{b_1, b_2, \dots, b_k\}$ for $n \geq 0$. As a main result, we estimate of their supports. The tools come from additive combinatorics and the uncertainty inequality.

- (15) **Russell Jay Hendel**, Towson University
Email: RHendel@Towson.Edu
Title: A system of four simultaneous recursions
Abstract: This paper further generalizes a recent result of Shannon and Ollerton who resurrected an old identity due to Ledin. This paper generalizes the Ledin-Shannon-Ollerton result to all metallic sequences. The results give closed formulas for the sum of products of powers of the first n integers with the first n members of the metallic sequence. Three key innovations of this paper are (i) reducing the proof of the generalization to the solution of a system of 4 simultaneous recursions; (ii) skillful use of the shift operation to prove equality of polynomials; and (iii) new OEIS sequences arising from the coefficients of the four polynomial families satisfying the 4 simultaneous recursions.
- (16) **Robert Hough**, Stony Brook University
Email: robert.hough@stonybrook.edu
Title: Character ratios and mixing on the symmetric and orthogonal groups
Abstract: I will describe new integral formulae for the characters of the symmetric and orthogonal groups, which give asymptotic evaluations of the characters. These character ratios have applications to several statistical physics models, including the Kac random walk on the orthogonal group, which models the velocities under collisions of N moving particles, and the random k -cycle walk on the symmetric group.
- (17) **Brad Isaacson**, New York City College of Technology (CUNY)
Email: bisaacson@citytech.cuny.edu
Title: On a polynomial reciprocity theorem of Carlitz
Abstract: Carlitz proved a powerful reciprocity theorem for generalized Dedekind-Rademacher sums. Among its many consequences was an interesting polynomial reciprocity theorem which holds under a certain restriction of its parameters. Carlitz remarked that it was unclear how this restriction could be removed. In this talk, we remove this restriction and obtain a generalization of Carlitz's polynomial reciprocity theorem.
- (18) **Faye Jackson**, University of Michigan, and
Luke Reifenberg, University of Notre Dame
Email: alephnil@umich.edu and lreifenb@nd.edu
Title: The Generalized Bergman game
Abstract: P. Baird-Smith A. Epstein, K. Flint, and S. J. Miler (2018) created the *Zeckendorf Game*, a two-player game which takes as an input a positive integer n and, using moves related to the Fibonacci recurrence relation, outputs the unique decomposition of n into a sum of non-consecutive Fibonacci numbers. Following this work and that of G. Bergman (1957), which proved the existence and uniqueness of such φ -decompositions, we formulate the *Bergman Game* which outputs the unique decomposition of n into a sum of non-consecutive powers of φ , the golden mean.

We then formulate *Generalized Bergman Games*, which use moves based on an arbitrary non-increasing positive linear recurrence relation and output the unique decomposition of n into a sum of non-adjacent powers of β , where β is the dominating root of the characteristic polynomial of the chosen recurrence relation. We prove that the longest possible Generalized Bergman game on an initial state S with n summands terminates in $\Theta(n^2)$ time, and we also prove that the shortest possible Generalized Bergman game on an initial state terminates between $\Omega(n)$ and $O(n^2)$ time. We also show a linear bound on the maximum length of the tuple used throughout the game.

This is joint work with Benjamin Baily, Justine Dell, Irfan Durmic, Henry Fleischmann, Isaac Mijares, Steven J. Miller, Ethan Pesikoff, Alicia Smith Reina, and Yingzi Yang.

- (19) **Renling Jin**, College of Charleston
 Email: jinr@cofc.edu
 Title: Hyper-hyper-hyper-integers and a simple proof of Szemerédi's theorem
 Abstract: In a conference five years ago, T. Tao reported his effort to simplify Szemerédi's original combinatorial proof of Szemerédi's theorem using nonstandard analysis. We continued his effort and presented a simple proof of the theorem for $k = 4$ in CANT 2020. In this talk, we will present a simple proof of the theorem for all k . One of the main simplifications is that a Tower of Hanoi type induction used by Szemerédi as well as Tao is replaced by a straightforward induction. In the proof the integers with three levels of infinities are used.
- (20) **Gergely Kiss**, Alfréd Rényi Institute of Mathematics, Hungary
 Email: kigergo57@gmail.com
 Title: Fuglede's conjecture on the direct product of finite abelian groups
 Abstract: We investigate Fuglede's conjecture on the direct product of abelian groups and its connection to the conjecture on \mathbb{R}^n for $n \geq 2$. We overview the earlier results: Some important constructions will be shown, which disproves the conjecture in higher dimensions, and some techniques and ideas will be presented, which serves to prove the conjecture for certain abelian groups. Finally we will discuss some developments of the most recent directions of research. This talk is closely related to Gábor Somlai's talk about Fuglede's conjecture on the cyclic group and one dimensional cases, and to the talk of Thomas Fallon who will present the detailed proof of Fuglede's conjecture for $\mathbb{Z}_p^2 \times \mathbb{Z}_q^2$.
- (21) **Jakub Konieczny**, Claude Bernard University Lyon 1, France
 Email: jakub.konieczny@gmail.com
 Title: Automatic semigroups
 Abstract: Automatic sequences, that is, sequences computable by finite automata, have been extensively studied from a variety of perspectives, including combinatorics, number theory, dynamics and theoretical computer science. Classification problems are a natural class of questions in

the theory of automatic sequences. In particular, the problem of classifying automatic multiplicative sequences has attracted considerable attention, culminating in complete classification which we obtained in joint work with Clemens Müllner and Mariusz Lemańczyk. The subject of my talk will be an extension of this line of inquiry, which we pursue in joint work with Oleksiy Klurman. Under mild technical assumptions, we classify all automatic multiplicative semigroups, that is, all sets E of integers which are closed under multiplication and such that the indicator function 1_E is automatic. Additionally, we show (again, under mild technical assumptions) that if E, F are automatic sets with $E \cdot F \subset E$ then E must contain a large essentially periodic component. This leads to potentially interesting open problems concerning products of automatic sets.

- (22) **Noah Kravitz**, Princeton University

Email: nkravitz@princeton.edu

Title: Zero patterns of derivatives of polynomials

Abstract: Motivated by recent work of Nathanson, we study the zero patterns of derivatives of polynomials. For P a polynomial of degree n and $\Lambda = (\lambda_1, \dots, \lambda_m)$ an m -tuple of distinct complex numbers, we consider the $m \times (n+1)$ *dope matrix* $D_P(\Lambda)$ whose ij -entry equals 1 if $P^{(j)}(\lambda_i) = 0$ and equals 0 otherwise (for $1 \leq i \leq m, 0 \leq j \leq n$). We address several natural questions: When m is 1 or 2, what do the possible dope matrices look like, and how many are there? What can we say about general upper bounds on the number of $m \times (n+1)$ dope matrices? For which m -tuples Λ is the number of $m \times (n+1)$ dope matrices maximized? Does every $\{0, 1\}$ -matrix appear as the left-most portion of some dope matrix?

Based on joint work with Noga Alon and Kevin O’Byrant.

- (23) **Thai Hoang Le**, University of Mississippi

Email: leth@olemiss.edu

Title: tba

- (24) **Paolo Leonetti**, Università “Luigi Bocconi”, Milano, Italy

Email: leonettipaolo@gmail.com

Title: The G.C.D. of n and the n th Fibonacci number

Abstract: Let $(F_n)_{n \geq 1}$ be the sequence of Fibonacci numbers, defined as usual by $F_1 = F_2 = 1$ and $F_{n+2} = F_{n+1} + F_n$ for all positive integers n ; and let \mathcal{A} be the set of all integers of the form $\gcd(n, F_n)$, for some positive integer n . In this talk we shall illustrate the following result on \mathcal{A} .

Theorem. *For all $x \geq 2$, we have*

$$\#\mathcal{A}(x) \gg \frac{x}{\log x}.$$

On the other hand, \mathcal{A} has zero asymptotic density.

The proofs rely on a result of Cubre and Rouse (PAMS, 2014) which gives, for each positive integer n , an explicit formula for the density of

primes p such that n divides the rank of appearance of p , that is, the smallest positive integer k such that p divides F_k .

- (25) **Jared Duker Lichtman**, University of Oxford
Email: jared.lichtman@maths.ox.ac.uk
Title: A proof of the Erdős primitive set conjecture
Abstract: A set of integers greater than 1 is primitive if no member in the set divides another. Erdős proved in 1935 that the series of $1/(n \log n)$, ranging over n in A , is uniformly bounded over all choices of primitive sets A . In 1988 he asked if this bound is attained for the set of prime numbers. In this talk we describe recent work which answers Erdős' conjecture in the affirmative. We will also discuss applications to old questions of Erdős, Sárközy, and Szemerédi from the 1960s.
- (26) **Ariane Masuda**, New York City College of Technology, CUNY
Email: amasuda@citytech.cuny.edu
Title: Rédei permutations with the same cycle structure
Abstract: Permutation polynomials over finite fields have been extensively studied over the past decades. Among the major challenges in this area are the questions concerning their cycle structures as they capture relevant properties, both theoretically and practically. In this talk we focus on a family of permutation polynomials, the so called Rédei permutations. Although their cycle structures are known, there are other related questions that can be investigated. For example, when do two Rédei permutations have the same cycle structure? We give a characterization of such pairs, and present explicit families of Rédei permutations with the same cycle structure. We also discuss some results regarding Rédei permutations with a particularly simple cycle structure, consisting of 1- and j -cycles only, when j is 4 or a prime number. The case $j = 2$ is specially important in some applications. We completely describe Rédei involutions with a prescribed cycle structure, and show that the only Rédei permutations with a unique cycle structure are the involutions.
This is joint work with Juliane Capaverde and Virgínia Rodrigues.
- (27) **Piotr Miska**, Jagiellonian University, Kraków, Poland
Email: piotr.miska@uj.edu.pl
Title: On (non-)realizability of Stirling numbers
Abstract: We say that a sequence $(a_n)_{n \in \mathbb{N}_+}$ of non-negative integers is realizable if there exists a set X and a mapping $T : X \rightarrow X$ such that a_n is the number of fixed points of T^n . For each $k \in \mathbb{N}_+$ and $j \in \{1, 2\}$ we define a sequence $S_k^{(j)} = (S^{(j)}(n + k - 1, k))_{n \in \mathbb{N}_+}$, where $S^{(j)}(n, k)$ is the Stirling number of the j -th kind (in case of $j = 1$ we consider unsigned Stirling numbers). The aim of the talk is to prove that $S_k^{(2)}$ is realizable if and only if $k \in \{1, 2\}$, while for $k \geq 3$ the sequence $S_k^{(2)}$ is almost realizable with a failure $(k - 1)!$, i. e. $(k - 1)!S_k^{(2)}$ is realizable. Moreover, I will show that for each $k \in \mathbb{N}_+$ the sequence $S_k^{(1)}$ is not almost realizable, i. e. for any $r \in \mathbb{N}_+$ the sequence $rS_k^{(1)}$ is not realizable.

The talk is based on a joint work with Tom Ward (Newcastle, UK).

- (28) **Mel Nathanson**, Lehman College (CUNY)
Email: melvyn.nathanson@lehman.cuny.edu
Title: Multiplicity interpolation of polynomials
Abstract: Interpolation problems related to the theorems of Descartes, Budan-Fourier, and Sturm in the theory of equations.
- (29) **Huy Pham**, Stanford University,
Email: huypham@stanford.edu
Title: Homogeneous structures in subset sums and applications
Abstract: In recent joint works with David Conlon and Jacob Fox, we develop novel techniques which allow us to prove a diverse range of results relating to subset sums. In the one-dimensional case, our techniques imply the existence of long *homogeneous arithmetic progressions* in the set of subset sums under a variety of assumptions. This allows us to resolve a number of longstanding open problems, including: solutions to the three problems of Burr and Erdős on Ramsey complete sequences, for which Erdős later offered a combined total of \$350; analogous results for the new notion of density complete sequences; the solution to a conjecture of Alon and Erdős on the minimum number of colors needed to color the positive integers less than n so that n cannot be written as a monochromatic sum; the exact determination of an extremal function introduced by Erdős and Graham on sets of integers avoiding a given subset sum; and, answering a question reiterated by several authors, a homogeneous strengthening of a result of Szemerédi and Vu on long arithmetic progressions in subset sums. In follow-up work in the multi-dimensional case, we show the existence of large *homogeneous generalized arithmetic progressions* in the set of subset sums of sufficiently large subsets of $[n]$, yielding a strengthening of a seminal result of Szemerédi and Vu. As an application, we make progress on the Erdős–Straus non-averaging sets problem, showing that every subset A of $[n]$ of size at least $n^{\sqrt{2}-1+o(1)}$ contains an element which is the average of two or more other elements of A . This gives the first polynomial improvement on a result of Erdős and Sárközy from 1990.
- (30) **Sean Prendiville**, Lancaster University, UK
Email: s.prendiville@lancaster.ac.uk
Title: Adapting the circle method for colourings
Abstract: Fix your favourite Diophantine equation. If each integer is coloured red, blue or green, how many solutions to your equation have all variables the same colour? We discuss how to adapt the Hardy-Littlewood circle method to yield a lower bound in certain problems of this flavour.
- (31) **Sinai Robins**, University of Sao Paulo, Brazil
Email: srobins@ime.usp.br
Title: The covariogram and an extension of Siegel’s formula
Abstract: We extend a formula of Carl Ludwig Siegel in the geometry of numbers. Siegel’s original formula assumed that there is exactly one lattice

point in the interior of the body, while here we relax that condition, so that the body may contain an arbitrary number of interior lattice points. Our extension involves a lattice sum of the covariogram for any compact set $\mathcal{K} \subset \mathbb{R}^d$, where the covariogram of \mathcal{K} at $x \in \mathbb{R}^d$ is defined by $\text{vol}(\mathcal{K} \cap (\mathcal{K} + x))$. The proof hinges on a variation of the Poisson summation formula which we derive here, and the Fourier methods herein also allow for more general admissible sets. One of the consequences of these results is a new characterization of multi-tilings of Euclidean space by translations, using the lower bound on lattice sums of such covariograms. The classical result known as Van der Corput's inequality, also follows immediately from the main result, as well as a new spectral formula for the volume of a compact set.

This is joint work with Michel Faleiros Martins.

- (32) **Carlo Sanna**, Politecnico di Torino, Italy

Email: carlo.sanna@polito.it

Title: Membership in random ratio sets

Abstract: Let \mathcal{A} be a random set constructed by picking independently each element of $\{1, \dots, n\}$ with probability $\alpha \in (0, 1)$. Several authors studied combinatorial/number-theoretic objects involving \mathcal{A} , including the sum set $\mathcal{A} + \mathcal{A}$, the product set $\mathcal{A}\mathcal{A}$, and the ratio set \mathcal{A}/\mathcal{A} . Generalizing a previous result of Cilleruelo and Gujarro-Ordóñez, we give a formula for the probability that a rational number q belongs to the ratio set \mathcal{A}/\mathcal{A} . Moreover, we give some results about formulas for the probability of the event $\bigvee_{i=1}^k (q_i \in \mathcal{A}/\mathcal{A})$, where q_1, \dots, q_k are rational numbers, showing that they are related to the study of the connected components of certain graphs. Finally, we provide some open question for future research.

- (33) **John R. Schmitt**, Middlebury College and University of Graz, Austria

Email: jschmitt@middlebury.edu

Title: Higher degree Davenport constants over finite commutative rings

Abstract: We generalize the notion of Davenport constants to a “higher degree” and obtain various lower and upper bounds, which are sometimes exact as is the case for certain finite commutative rings of prime power cardinality. Two simple examples that capture the essence of these higher degree Davenport constants are the following. 1) Suppose $n = 2^k$, then every sequence of integers S of length $2n$ contains a subsequence S' of length at least two such that $\sum_{a_i, a_j \in S'} a_i a_j \equiv 0 \pmod{n}$ and the bound is sharp. 2) Suppose $n \equiv 1 \pmod{2}$, then every sequence of integers S of length $2n - 1$ contains a subsequence S' of length at least two such that $\sum_{a_i, a_j \in S'} a_i a_j \equiv 0 \pmod{n}$. These examples illustrate that if a sequence of elements from a finite commutative ring is long enough, certain symmetric expressions have to vanish on the elements of a subsequence.

Joint work with Yair Caro and Benjamin Girard.

- (34) **James Sellers**, University of Minnesota Duluth
 Email: jsellers@d.umn.edu
 Title: Relating the crank of a partition and smallest missing parts
 Abstract: The primary goal of this talk is to demonstrate a natural connection between the smallest missing part of an integer partition (commonly referred to as the “mex” of the partition) and the concept of the crank of a partition. After providing a brief history of the crank of a partition a la Dyson as well as Andrews and Garvan, we will utilize straightforward generating function manipulations to make this connection. We will then consider additional results on the mex statistic based on parity, and we will also demonstrate connections between the crank and Frobenius symbols which satisfy certain conditions.
 This work is joint with Brian Hopkins, Dennis Stanton, and Ae Ja Yee.
- (35) **Steven Senger**, Missouri State University
 Email: stevensenger@gmail.com
 Title: Distinct dot products, convexity, and $AA + 1$
 Abstract: We discuss recent developments in estimating the number of distinct dot products determined by a large finite set of n points in the plane. The improvement comes from improved understanding of the multiplicative structure of an additively shifted product set, $AA + 1$, when A is a large finite subset of the real numbers. This breakthrough was made possible by new additive combinatorial results about convex sets of numbers.
- (36) **Junxuan Shen**, California Institute of Technology
 Email: jshen@caltech.edu
 Title: The structural incidence problem for cartesian products
 Abstract: We prove new structural results for point-line incidences. An incidence is a pair of one point and one line, where the point is on the line. The Szemerédi-Trotter theorem states that n points and n lines form $O(n^{4/3})$ incidences. This bound has been used to obtain many results in combinatorics, number theory, harmonic analysis, and more. While the Szemerédi-Trotter bound has been known for several decades, the structural problem remains wide-open. This problem asks to characterize the point-line configurations with $\Theta(n^{4/3})$ incidences. We prove that when the point set \mathcal{P} is a Cartesian product where only one axis of it behaves like a lattice, the line set must contain many families of parallel lines to achieve the maximal incidence bound.
 Theorem: Consider $1/3 < \alpha < 2/3$. Let $A, B \subset {}^*\mathbf{R}$ satisfy that $A = \{1, 2, \dots, n^\alpha\}$ and $|B| = n^{1-\alpha}$. Let \mathcal{L} be a set of n lines in ${}^*\mathbf{R}^2$, such that $I(A \times B, \mathcal{L}) = \Theta(n^{4/3})$. Then \mathcal{L} contains $\Omega(n^{1-\beta}/\log n)$ disjoint families of $\Theta(n^\beta)$ parallel lines for $1 - 2\alpha \leq \beta \leq 2/3$.
 When $\alpha < 1/3$ or $\alpha > 2/3$, it is impossible to have $\Theta(n^{4/3})$ incidences. We also completely characterize the line set when the point set is a lattice.
 Joint work with Adam Sheffer.

- (37) **Bartosz Sobolewski**, Jagiellonian University, Kraków, Poland
 Email: bartosz.sobolewski@uj.edu.pl
 Title: Monochromatic arithmetic progressions in binary words associated with pattern sequences
 Abstract: Let $e_v(n)$ denote the number of occurrences of a pattern v in the binary expansion of $n \in \mathbb{N}$. In the talk we consider monochromatic arithmetic progressions in the class of words $(e_v(n) \bmod 2)_{n \geq 0}$ over $\{0, 1\}$, which includes the Thue–Morse word \mathbf{t} ($v = 1$) and a variant of the Rudin–Shapiro word \mathbf{r} ($v = 11$). So far, the problem of exhibiting long progressions and finding an upper bound on their length has mostly been studied for \mathbf{t} and certain generalizations. We show that analogous results hold for \mathbf{r} . In particular, we prove that a monochromatic arithmetic progression of difference $d \geq 3$ starting at 0 in \mathbf{r} has length at most $(d + 3)/2$, with equality infinitely often. We also compute the maximal length of progressions of differences $2^k - 1$ and $2^k + 1$. Some weaker results for a general pattern v are provided as well.
- (38) **Gábor Somlai**, Eötvös Loránd University and Alfréd Rényi Institute of Mathematics, Hungary
 Email: zsomlai@gmail.com
 Title: Fuglede’s conjecture, the one dimensional case
 Abstract: Fuglede conjectured that a bounded measurable set (in \mathbb{R}^n) is spectral if and only if it is a tile. The conjecture was also confirmed by Fuglede for sets whose tiling complement is lattice and for spectral sets one of whose spectrums is a lattice. The conjecture was disproved by Tao by constructing a spectral set in \mathbb{Z}_3^5 , which is not a tile and lifted it to the 5 dimensional Euclidean space.
 The conjecture is open only in dimensions 1 and 2. The 1 dimensional case is directly connected with the one of finite cyclic groups and to the so called Coven-Meyerowitz conjecture. One of the main aims of the talk is to present some of the methods developed that lead to our recent results.
- (39) **Johann Thiel**, New York City College of Technology, CUNY
 Email: jthiel@citytech.cuny.edu
 Title: Solving the membership problem for certain subgroups of $SL_2(\mathbb{Z})$
 Abstract: For positive integers u and v , let $L_u = \begin{bmatrix} 1 & 0 \\ u & 1 \end{bmatrix}$ and $R_v = \begin{bmatrix} 1 & v \\ 0 & 1 \end{bmatrix}$. Let $G_{u,v}$ be the group generated by L_u and R_v . The membership problem for $G_{u,v}$ asks the following question: Given a 2-by-2 matrix $M = \begin{bmatrix} a & b \\ c & d \end{bmatrix}$, is there a relatively straightforward method for determining if M is a member of $G_{u,v}$? In the case where $u = 2$ and $v = 2$, Sanov was able to show that simply checking some divisibility conditions for a , b , c and d is enough to make this determination. We answered this question in the case where $u, v \geq 3$ by finding a characterization of matrices M in $G_{u,v}$ in terms of the short continued fraction representation of $\frac{b}{a}$, extending some results of Esbelin and Gutan. By modifying our previous work, we are able to further

extend our previous result to the more difficult case where $u, v \geq 2$ with $uv \neq 4$.

This is joint work with Sandie Han, Ariane M. Masuda, and Satyanand Singh.

- (40) **Ognian Trifonov**, University of South Carolina

Email: trifonov@math.sc.edu

Title: Lattice points close to ovals, arcs, and helices

Abstract: In 1972 Schinzel showed that the largest distance between three lattice points on a circle of radius R is at least $\sqrt[3]{2}R^{1/3}$. We generalize Schinzel's result to ovals and arcs with bounded curvature in the plane and lattice points close to the curve. Furthermore, we extend the result to the case of affine lattices. Finally, we obtain similar results when the curve is a helix in three dimensional space.

- (41) **Maciej Ulas**, Jagiellonian University, Kraków, Poland

Email: maciej.ulas@gmail.com

Title: Solutions of certain meta-Fibonacci recurrences

Abstract: We investigate the solutions of certain meta-Fibonacci recurrences of the form $f(n) = f(n - f(n - 1)) + f(n - 2)$ for various sets of initial conditions. In the case when $f(n) = 1$ for $n \leq 1$, we prove that the resulting integer sequence is closely related to the function counting binary partitions of a certain type (independently of the value of $f(2) \in \mathbb{N}$).

The talk is based on a joint work with Bartosz Sobolewski.

- (42) **Ethan Patrick White**, University of British Columbia

Email: epwhite@math.ubc.ca

Title: Erdős' minimum overlap problem

Abstract: In 1955 Erdős posed the following problem. Let n be a positive integer and $A, B \subset [2n]$ be a partition of $[2n]$ such that $|A| = |B| = n$. For any such partition and integer $-2n < k < 2n$, define M_k to be the number of solutions $(a, b) \in A \times B$ to $a - b = k$. Estimate the size of the function

$$M(n) = \min_{A \cup B = [2n]} \max_{-2n < k < 2n} M_k,$$

where the minimum is taken over all partitions of $[2n]$ into equal-sized sets. Many upper and lower estimates were obtained over the following decades, and the state of the art is $0.356 < M(n)/n < 0.381$. We use elementary Fourier analysis to translate the problem to a convex optimization program and obtain the new lower bound $M(n)/n > 0.379$.

- (43) **Trevor D. Wooley**, Purdue University
 Email: twooley@purdue.edu
 Title: Shifted analogues of the divisor function
 Abstract: Suppose that θ is irrational. Then almost all elements $\nu \in \mathbb{Z}[\theta]$ that may be written as a k -fold product of the shifted integers $n + \theta$ ($n \in \mathbb{N}$) are thus represented essentially uniquely. We discuss this and related paucity problems.
 Most of this work is joint with Winston Heap and Anurag Sahay.
- (44) **Max Wenqiang Xu**, Stanford University
 Email: maxxu@stanford.edu
 Title: On a Turán Conjecture and random multiplicative functions
 Abstract: We show that if f is the random completely multiplicative function, the probability that $\sum_{n \leq x} \frac{f(n)}{n}$ is positive for every x is at least $1 - 10^{-40}$. For large x we prove an asymptotic upper bound of $O(\exp(-\exp(\frac{\log x}{C \log \log x})))$ on the probability that a particular truncation is negative.
 This is joint work with Rodrigo Angelo.
- (45) **Catherine Yan**, Texas A&M University
 Email: huafei-yan@tamu.edu
 Title: Multivariate Gončarov polynomials and integer sequences
 Abstract: Univariate delta Gončarov polynomials arise when the classical Gončarov interpolation problem in numerical analysis is modified by replacing derivatives with delta operators. When the delta operator under consideration is the backward difference operator, we acquire the univariate difference Gončarov polynomials, which have a combinatorial relation to lattice paths in the plane with a given right boundary. In this talk, we extend several algebraic and analytic properties of univariate Gončarov polynomials to the multivariate case with both the derivative and backward difference operators. We then establish a combinatorial interpretation of multivariate Gončarov polynomials in terms of certain constraints on d -tuples of integer sequences. This motivates a connection between multivariate Gončarov polynomials and a higher-dimensional generalized parking function, the **U**-parking function, from which we derive several enumerative results based on the theory of delta operators.
 This talk is based on joint work with Ayo Adeniran and Lauren Snider.
- (46) **Daodao Yang**, Graz University of Technology, Austria
 Email: yang@tugraz.at
 Title: Extreme values of derivatives of the Riemann zeta function, log-type GCD sums, and spectral norms
 Abstract: First I will recall the research on greatest common divisor (GCD) sums and extreme values of the Riemann zeta function. The motivation for the study and the connection between the two problems will be discussed. Then I will explain how to establish lower bounds for maximums of $|\zeta^{(\ell)}(\sigma + it)|$ when $\sigma \in [\frac{1}{2}, 1]$, $\ell \in \mathbb{N}$. One of my results states that as

$T \rightarrow \infty$, uniformly for all positive integers $\ell \leq (\log_3 T)/(\log_4 T)$, we have

$$\max_{T \leq t \leq 2T} \left| \zeta^{(\ell)}(1+it) \right| \geq (\mathbf{Y}_\ell + o(1)) (\log_2 T)^{\ell+1},$$

where $\mathbf{Y}_\ell = \int_0^\infty u^\ell \rho(u) du$, and $\rho(u)$ denotes the Dickman function. This generalizes results of Bohr-Landau and Littlewood on $|\zeta(1+it)|$ in 1910s. The tools are Soundararajan's resonance methods and ingredients are certain combinatorial optimization problems. On the other hand, assuming the Riemann hypothesis, we have $|\zeta^{(\ell)}(1+it)| \ll_\ell (\log \log t)^{\ell+1}$. Then I will talk on the log-type GCD sums $\Gamma_\sigma^{(\ell)}(N)$, which I define it as follows

$$\Gamma_\sigma^{(\ell)}(N) := \sup_{|\mathcal{M}|=N} \frac{1}{N} \sum_{m,n \in \mathcal{M}} \frac{(m,n)^\sigma}{[m,n]^\sigma} \log^\ell \left(\frac{m}{(m,n)} \right) \log^\ell \left(\frac{n}{(m,n)} \right),$$

where the supremum is taken over all subsets $\mathcal{M} \subset \mathbb{N}$ with size N . I will explain how $\Gamma_\sigma^{(\ell)}(N)$ can be related to $|\zeta^{(\ell)}(1+it)|$ and how to prove that $(\log \log N)^{2+2\ell} \ll_\ell \Gamma_1^{(\ell)}(N) \ll_\ell (\log \log N)^{2+2\ell}$, which generalizes Gál's theorem (corresponding to the case $\ell = 0$). The lower bounds could be used to produce large values of $|\zeta^{(\ell)}(1+it)|$. Using a random model for the zeta function via methods of Lewko-Radziwiłł, upper bounds for spectral norms on α -line are established, when $\alpha \rightarrow 1^-$ with certain fast rates. As a corollary, upper bounds of correct order of the log-type GCD sums are established.

(47) **Wijit Yangjit**, University of Michigan

Email: yangjit@umich.edu

Title: On the Montgomery-Vaughan weighted generalization of Hilbert's inequality

Abstract: Hilbert's inequality states that

$$\left| \sum_{m=1}^N \sum_{\substack{n=1 \\ n \neq m}}^N \frac{z_m \bar{z}_n}{m-n} \right| \leq C_0 \sum_{n=1}^N |z_n|^2,$$

where C_0 is an absolute constant. In 1911, Schur showed that the optimal value of C_0 is π .

In 1974, Montgomery and Vaughan proved a weighted generalization of Hilbert's inequality and used it to estimate mean values of Dirichlet series. This generalized Hilbert inequality is important in the theory of the large sieve. The optimal constant C in this inequality is known to satisfy $\pi \leq C < \pi + 1$. It is widely conjectured that $C = \pi$. In this talk, I will describe the known approaches to obtain an upper bound for C , which proceed via a special case of a parametric family of inequalities. We analyze the optimal constants in this family of inequalities. A corollary is that the method in its current form cannot imply an upper bound for C below 3.19.

- (48) **Chi Hoi Yip**, University of British Columbia
Email: kyleyip@math.ubc.ca
Title: Asymptotics for the number of directions determined by $[n] \times [n]$ in \mathbb{F}_p^2
Abstract: Let p be a prime and n a positive integer such that $\sqrt{\frac{p}{2}} + 1 \leq n \leq \sqrt{p}$. For any arithmetic progression A of length n in \mathbb{F}_p , we establish an asymptotic formula for the number of directions determined by $A \times A \subset \mathbb{F}_p^2$. The key idea is to reduce the problem to counting the number of solutions to the bilinear Diophantine equation $ad + bc = p$ in variables $1 \leq a, b, c, d \leq n$; our asymptotic formula for the number of solutions is of independent interest.
Joint work with Greg Martin and Ethan White.
- (49) **Qinghai Zhong**, University of Graz, Austria,
Email: qinghai.zhong@uni-graz.at
Title: On monoids of weighted zero-sum sequences
Abstract: Let G be an additive finite abelian group and $\Gamma \subset \text{End}(G)$ be a subset of the endomorphism group of G . A sequence $S = g_1 \cdot \dots \cdot g_\ell$ over G is a (Γ) -weighted zero-sum sequence if there are $\gamma_1, \dots, \gamma_\ell \in \Gamma$ such that $\gamma_1(g_1) + \dots + \gamma_\ell(g_\ell) = 0$. We study algebraic and arithmetic properties of monoids of weighted zero-sum sequences.